# **Gastrointestinal Associates LLP**

210 Portland Street, Suite 100 Columbia, MO 65201 573-777-8820

Dr. Peter Cleavinger, Dr. Michele Diaz-Arias, Dr. Donald Gerhardt, Dr. Kevin Halsey, Dr. Nicolas Llorens, Dr. Matthew Struttmann, and Dr. Stephen Welsh

Your procedure is sched	<u>luled:</u>	
[ ] Columbia Endoscopy	Center (573)449-3500, 20	08 Portland St., Columbia, MO 65201
Date:	Arrive at:	Procedure Time:
[ ] [ ] [ ] [ ] [ [ ] [ ] [ ]	145) 0000 4000 F 4 B	
	-	adway, Columbia, MO 65201
Date:	Arrive at:	Procedure Time:
	MoviPrep Bo	wel Prep
past 30 years, it has beco- colorectal cancer. With m instruments, most exams before. For an accurate a material by means of a re	me a gold standard in scr lodern methods of sedation are quick, painless, and v and safe exam, it is very in stricted diet and laxative p	examine the colon or large intestine. Over the reening for colon polyps and prevention of on and newer computerized flexible very safe. The major obstacle is the day important that your colon be clean of all waste preparation or "prep". Follow all instructions luled. If you have any questions please call
To prepare for your cold	noscopy you will need	to purchase:
[X] One <b>MOVIPREP</b> Bo	wel Prep Kit (you will need	d a prescription)
All bowel preps may caus	e some anal irritation. We	e also recommend:
[ ] A soft brand of toilet ti	ssue (such as Charmin Se	ensitive)
[ ] A box of baby wipes o	r tucks pads (such as Huເ	ggies or Pampers Aloe wipes)
[ ] A small tube of diaper	rash or Zinc Oxide cream	ı (such as A&D Zinc Oxide)
		,

### When your appointment is scheduled:

If you are on any **blood thinning medications/anticoagulants** (Coumadin, Jantoven, Warfarin, Plavix, Pradaxa, Effient, Xarelto, Brilinta, Aggrenox, Ticlid, Aggrastat, Avixtra, Iprivask, Lovenox, Dipyridamole), please contact our office at 573-777-8820. These medications often need to be stopped 5 days prior to the procedure.

### Three to Five days before the your colonoscopy:

- 1) Make arrangements for a responsible adult to accompany you to your appointment and drive you home. If possible, this person should be someone we can share the results of the exam with.
- 2) Stop taking medications that contain iron.
- 3) Continue any other medications that you normally take.
- 4) For females, if you have started your menstrual period it is still okay to do the colonoscopy. Female patients of the appropriate age will be asked to provide a urine specimen for a pregnancy test the morning of your procedure at the facility.
- 5) It is helpful to eat lightly a few days before your exam. It makes the cleanout easier and more effective. Try to avoid overeating to compensate for the upcoming fast. Please avoid Metamucil, raw vegetables, corn, and foods with skin or small seeds (like tomatoes, sesame seeds, kiwi, and cucumbers) for a few days before your exam.

#### The day before your colonoscopy:

- 1) Eat a clear liquid diet for breakfast, lunch, and dinner the day before the test. CLEAR LIQUIDS are anything you can see through. This includes clear water, fat free broths, bouillon, tea, coffee (without creamer), Kool-Aid, Crystal Light, carbonated beverages, sodas, fruit juices without pulp (apple, white grape, cranberry), popsicles without pulp, Jell-O, hard candies, Italian ices, Gatorade. AVOID anything with red dye. NOT ALLOWED: Orange juice, fruit nectars, liquids you cannot see through, milk, cream. Please drink plenty of fluids during the day to prevent dehydration.
- 2) If you are a diabetic patient on insulin, please check with your **primary physician** regarding necessary changes to your doses.
- 3) Between 12:00p- 5:00PM, mix the contents of 1-pouch A and 1-pouch B with 1-liter of water in the mixing container. The MoviPrep container is divided by 4 lines. Every 15 minutes drink the solution down to the next line (about 8 ounces) until the container is empty. For a safe prep and complete exam you MUST drink another 32 ounces of clear fluid over the next hour. You can mix the solution ahead of time and refrigerate it prior to drinking—solution should be used within 24 hours of mixing. Your stools should eventually be a clear yellow color and you should be able to see the bottom of the toilet through the fluid after prepping.
- 4) You may continue drinking clear liquids throughout the evening.

## The day of your colonoscopy:

- 1) No breakfast
- 2) At least 4 hours before your appointment, mix the contents of 1-pouch A and 1-pouch B with 1-liter of water in the mixing container. The MoviPrep container is divided by 4 lines. Every 15 minutes drink the solution down to the next line (about 8 ounces) until the container is empty. For a safe prep and complete exam you MUST drink another 32 ounces of clear fluid over the next hour. You must finish drinking the final glass of water

- at least 3 hours before your check in time. You need to do the second dose the morning of your procedure even if your stools were clear the night before.
- 3) You cannot have any food, drink, candy, gum, chewing tobacco and should not smoke for 3 hours before your procedure. You should not have broth the morning of your procedure.
- 4) You may take blood pressure medications, seizure medications, prophylactic antibiotics, and inhalers prior to your procedure with a sip of water. Take insulin as directed by your primary physician.
- 5) Dress comfortably. You may want to bring a change of clothes just in case you soil the clothes you are wearing. You may want to consider wearing Depends if you have a long drive.
- 6) You will need to bring a driver, a list of your medications/vitamins (that includes the name, dose, frequency, and reason for taking the medication), driver's license (or Identification), insurance card, and co-pay if your insurance requires one. Leave your jewelry and other valuables at home.

### After arriving at the Endoscopy Center:

- 1) After arriving at the Center/Hospital, you are to check in. Sedatives/anesthesia will be used for the procedure to help with any discomfort you may have, therefore an IV will be placed in your arm/hand.
- 2) You will need to sign a consent form authorizing the physician to perform the procedure. The procedure takes 20-30 minutes, and recovery is about 30 minutes. Your total time at the Center/Hospital should be approximately 2-3 hours. Before you leave, you will receive information about the results of your colonoscopy, instructions regarding diet, IV, and follow up.
- 3) You will not be able to drive or return to work for the remainder of the day.

If you have any questions concerning the colonoscopy or prep, please do not hesitate to call our office at 573-777-8820.